



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE ON CAUCHY'S INTEGRAL TEST.

By M. B. PORTER, University of Texas.

It does not seem to have been generally noticed that series which consist of a monotonic decreasing sequence of terms of the same sign lend themselves to calculation as readily as those whose signs alternate provided the integral $\int_n^\infty u_n dn$ is sufficiently simple. We have at once, if the terms are all positive,

$$\int_n^\infty u_n dn \leq u_n + u_{n+1} + u_{n+2} + \dots \text{etc.} \leq u_n + \int_n^\infty u_n dn.$$

Thus if to the sum of the first $n-1$ terms we add $\frac{u_n}{2} + \int_n^\infty u_n dn$, our approximation will not differ from the sum of the series by more than $u_n/2$, that is, by less than half of the first term we omit.

Thus to calculate the series $\sum_1^\infty n^{-2}$ the correction after $n-1$ terms is $\frac{1}{2n^2} + \frac{1}{n}$, and the calculated value is in error not more than $\frac{1}{2n^2}$. This fact can be used to calculate with a minimum of labor such slowly convergent series as those represented by the logarithmic scales of DeMorgan.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

346. Proposed by E. B. ESCOTT, Professor of Mathematics, University of Michigan.

Solve completely by quadratics:

$$\frac{2x}{1-x^2}=y; \quad \frac{2y}{1-y^2}=z; \quad \frac{2z}{1-z^2}=w; \quad \frac{2w}{1-w^2}=x.$$

Solution by the PROPOSER.

Let $x=\tan\phi$, then $y=\tan 2\phi$, $z=\tan 4\phi$, $w=\tan 8\phi$, $x=\tan 16\phi=\tan\phi$,
 $\tan 16\phi - \tan\phi = 0$, or $\frac{\sin 15\phi}{\cos 16\phi \cdot \cos\phi} = 0$.